## Measurement of the Thermal Properties of Co(NO<sub>3</sub>)<sub>2</sub> and Acetamide by a Modified AC-Heated Wire Technique

S.R. Atalla, G.A. Attia, M.M. El-Naggar, and T.A. El-Sayed

Department of Physics

Faculty of Education

Fayoum Branch

Cairo University, Faycum, Egypt

A.A. EI-Sharkawy Department of Physics Faculty of Science Al-Azhar University Cairo, Egypt

## N. Kamel

Department of Physics Faculty of Science Ain Shams University Cairo, Egypt

M. Kenawy
Department of Physics
Faculty of Women
Ain Shams University
Cairo, Egypt

A modified AC-heated wire technique has been used for the measurement of thermal activity (b), thermal diffusivity (a). volumetric heat capacity ( $\rho C_p$ ) and the thermal conductivity ( $\lambda$ ) of nitrate salt  $Co(N0_3)_2$  in the temperature range  $30\text{-}120^{\circ}C$  and acetamide  $CH_3CONH_2$  in the temperature range  $50\text{-}90^{\circ}C$ . DSC investigations have been also performed for both substances.

The results for  $Co(N0_3)_2$  show that (a) slightly decreases,  $(\rho C_p)$  increases but  $\lambda$  remains constant in the measured temperature range. For acetamide, (a) slightly decreases,  $(\rho C_p)$  increases, and  $(\lambda)$  increases with temperature, but all of them do not change appreciably on melting. The obtained values for  $\lambda$  for both substances show that the main mechanism of heat transport is due to phonons as expected.

The upper limit of the investigated temperature ranges was dictated by the occurrence of unbalance of the measuring system and disappearance of the third harmonic signal. This observation requires further investigations to find out the reason behind it, especially because the same phenomenon has been observed for some other salts.